

• 第23讲: 重积分应用举例

例1. 设物体 $\Omega: \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} \leq 1$ 的体密度为 $\rho(x,y,z) =$

$x^2 + y^2 + z^2$, 求 Ω 的总质量 $M(\Omega)$

这也是 $\rho_0 = 1$ 的物体 Ω 关于原点的转动惯量

解: $M(\Omega) = \iiint_{\Omega} (x^2 + y^2 + z^2) dV = \iiint_{\Omega} x^2 dV + \iiint_{\Omega} y^2 dV + \iiint_{\Omega} z^2 dV$

• 而 $\iiint_{\Omega} x^2 dV = \int_{-a}^a x^2 dx \iint_{D_x} 1 dy dz$, $D_x: \frac{y^2}{b^2} + \frac{z^2}{c^2} \leq 1 - \frac{x^2}{a^2}$

$= \int_{-a}^a x^2 \cdot S(D_x) dx = \int_{-a}^a x^2 \cdot 2bc \sqrt{1 - \frac{x^2}{a^2}} dx = \frac{4}{15} \pi a^3 bc$

同理, $\iiint_{\Omega} y^2 dV = \frac{4}{15} \pi b^3 ac$, $\iiint_{\Omega} z^2 dV = \frac{4}{15} \pi c^3 ab$

$\therefore M(\Omega) = \frac{4}{15} \pi abc(a^2 + b^2 + c^2)$

• 例2. 求物体 $\Omega: \left(\frac{x}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{b}\right)^{\frac{2}{3}} + \left(\frac{z}{c}\right)^{\frac{2}{3}} \leq 1$ 的体积 $V(\Omega)$

解: $V(\Omega) = \iiint_{\Omega} 1 dV = \iiint_{\Omega} 1 dx dy dz$ $x = au^3, y = bv^3, z = cw^3$

$\iiint_{u^2+v^2+w^2 \leq 1} \left| \frac{\partial(x,y,z)}{\partial(u,v,w)} \right| du dv dw = \iiint_{u^2+v^2+w^2 \leq 1} 7abc u^2 v^2 w^2 du dv dw$

$u = r \sin \theta \cos \phi, v = r \sin \theta \sin \phi$

$w = r \cos \theta$

$= 7abc \int_0^{2\pi} d\phi \int_0^{\pi} \int_0^1 (r \sin \theta \cos \phi)^2 (r \sin \theta \sin \phi)^2 (r \cos \theta)^2 r^2 \sin \theta dr$

$$\begin{aligned}
 &= 27abc \int_0^{2\pi} \cos^2 \theta \sin^2 \theta d\theta \int_0^{\frac{\pi}{2}} \sin^5 \theta \cos^2 \theta d\theta \int_0^1 r^8 dr \\
 &= 27abc \times 4 \int_0^{\frac{\pi}{2}} (\cos^2 \theta \sin^2 \theta d\theta) \times 2 \int_0^{\frac{\pi}{2}} (\sin^5 \theta - \sin^7 \theta) d\theta \times \frac{1}{9} \\
 &= 12abc \left(\frac{1}{2} \times \frac{2}{2} - \frac{3}{4} \times \frac{1}{2} \times \frac{2}{2} \right) \times 2 \left(\frac{4}{5} \times \frac{2}{3} \times 1 - \frac{6}{7} \times \frac{4}{5} \times \frac{2}{3} \right) = \frac{4}{35} \pi.
 \end{aligned}$$

例3. 求 $\int_{\Omega} f(x, y, z) = x^2 + y^2 + z^2$ 在 $\Omega = x^2 + y^2 + z^2 \leq x + y + z$ 上的二重积分

积分平均值 \bar{f} .

$$\text{解: } \bar{f} = \frac{\iiint_{\Omega} f(x, y, z) dV}{V(\Omega)}.$$

而 $\Omega = (x - \frac{1}{2})^2 + (y - \frac{1}{2})^2 + (z - \frac{1}{2})^2 \leq (\frac{\sqrt{3}}{2})^2$ 为球 $V(\Omega) =$

$$\frac{4}{3} \pi R^3 = \frac{4}{3} \pi \left(\frac{\sqrt{3}}{2} \right)^3 = \frac{\sqrt{3}}{2} \pi. \text{ 且}$$

$$\iiint_{\Omega} f(x, y, z) dV = \int_0^{2\pi} \int_0^{2\pi} \int_0^{\frac{\sqrt{3}}{2}} \left[\left(\frac{1}{2} + r \sin \theta \cos \varphi \right)^2 + \left(\frac{1}{2} + r \sin \theta \sin \varphi \right)^2 + \left(\frac{1}{2} + r \cos \theta \right)^2 \right] r^2 \sin \theta dr d\theta d\varphi$$

$$= \int_0^{2\pi} \int_0^{2\pi} \int_0^{\frac{\sqrt{3}}{2}} \left[\frac{3}{4} r^2 \sin^2 \theta + r^2 \sin^2 \theta (\cos^2 \varphi + \sin^2 \varphi) + r^2 \cos^2 \theta + \frac{4}{2} r \sin \theta \cos \theta \right] r^2 \sin \theta dr d\theta d\varphi$$

$$= \int_0^{2\pi} \int_0^{2\pi} \int_0^{\frac{\sqrt{3}}{2}} \left[\frac{3}{4} r^2 \sin^2 \theta + r^2 \sin^2 \theta (\cos^2 \varphi + \sin^2 \varphi) + r^2 \cos^2 \theta + \frac{4}{2} r \sin \theta \cos \theta \right] r^2 \sin \theta dr d\theta d\varphi$$

$$= \int_0^{2\pi} \int_0^{2\pi} \int_0^{\frac{\sqrt{3}}{2}} \left(\frac{3}{4} r^2 \sin^2 \theta + r^2 \sin^2 \theta + r^2 \cos^2 \theta + 2r \sin \theta \cos \theta \right) r^2 \sin \theta dr d\theta d\varphi$$

$$= 2\pi \int_0^{2\pi} \int_0^{\frac{\sqrt{3}}{2}} \left(\frac{3}{4} r^2 + r^2 + r^2 \cos^2 \theta + 2r \sin \theta \cos \theta \right) r^2 \sin \theta dr d\theta = 2\pi \times 2 \left(\frac{1}{4} \times \frac{3\sqrt{3}}{8} + \frac{9\sqrt{3}}{5 \times 32} \right) = \frac{3}{5} \sqrt{3} \pi.$$

$$\therefore \bar{f} = \frac{3}{5} \sqrt{3} \pi / \frac{\sqrt{3}}{2} \pi = \frac{6}{5}. \quad (\text{解法(2)见后面}) \quad (2)$$

- 例4. 证明刚体转动惯量的平行轴定理:

$$J_k = J_c + m \cdot d^2 \quad (d \text{ 是两轴之间的距离})$$

m 为物体 Ω 的质量: $m = \iiint_{\Omega} \rho(x, y, z) dV$. J_c 是 Ω 绕


过物体重心 $G(x, y, z)$ 之轴 C 的转动惯量, J_k 是 Ω 绕

- 与轴 C 平行的轴 k 的转动惯量.

证 (1) 建立坐标系, 使轴 k

与 z 轴重合, 且重心 $G(x, y, z)$

位于 oy 轴上: $G(0, d, 0)$



- 即 $\bar{x} = \frac{\iiint_{\Omega} x \rho(x, y, z) dV}{m} = \bar{z} = \frac{\iiint_{\Omega} z \rho(x, y, z) dV}{m} = 0$,

$$\bar{y} = \frac{\iiint_{\Omega} y \rho(x, y, z) dV}{m} = d \Rightarrow \iiint_{\Omega} y \rho(x, y, z) dV = md.$$

- (2) $J_k = J_z = \iiint_{\Omega} (x^2 + y^2) \rho(x, y, z) dV$

$$J_c = \iiint_{\Omega} (x^2 + (y-d)^2 + 0^2) \rho(x, y, z) dV = J_k + \iiint_{\Omega} d^2 \rho dV - 2d \iiint_{\Omega} y \rho dV$$

$$= J_k + d^2 \cdot m - 2d(m \cdot d) = J_k - d \cdot m \Leftrightarrow J_k = J_c + m \cdot d^2$$

- 例5. 求半径为 R 的均匀球体 Ω 对球外一质点 Q 的引力 $\vec{F} = (F_x, F_y, F_z)$, 其中质点 Q 的质量为 m_0 .


解: (1) 设均匀球体 $\Omega: x^2 + y^2 + z^2 \leq R^2$ 的体密度为 $\rho_0 > 0$.

(2) 建立坐标系, 使 Q 点位于 z 轴上 $(0, 0, a)$ 处, ($a > R > 0$)

(3) 在 Ω 中任取一子区域 Ω_i , 质量 $m(\Omega_i) = \rho_0 dV$.

$$d\vec{F} = k \frac{m_0 (\rho_0 dV)}{x^2 + y^2 + (z-a)^2} \cdot \vec{F}_0$$

$$\vec{F}_0 = \frac{(x, y, z-a)}{\sqrt{x^2 + y^2 + (z-a)^2}}$$



$$dF_x = \frac{k m_0 \rho_0 x dx dy dz}{(x^2 + y^2 + (z-a)^2)^{3/2}}, \quad dF_y = \frac{k m_0 \rho_0 y dx dy dz}{(x^2 + y^2 + (z-a)^2)^{3/2}}$$

$$dF_z = \frac{k m_0 \rho_0 (z-a) dx dy dz}{(x^2 + y^2 + (z-a)^2)^{3/2}}, \quad \text{由对称性可知:}$$

$$\iiint_{\Omega} dF_x = F_x = 0 = F_y = \iiint_{\Omega} dF_y \quad \text{而} \quad F_z = \iiint_{\Omega} dF_z \quad \begin{matrix} x=r\cos\theta \\ y=r\sin\theta \\ z=z \end{matrix}$$

$$k m_0 \rho_0 \int_{-R}^R (z-a) dz \int_0^{2\pi} \int_0^{\sqrt{R^2 - z^2}} \frac{r dr}{(r^2 + (z-a)^2)^{3/2}} d\theta$$

$$= 2\pi k m_0 \rho_0 \int_{-R}^R (z-a) (r^2 + (z-a)^2)^{-1/2} \Big|_0^{\sqrt{R^2 - z^2}} dz$$

$$= -2\pi k m_0 \rho_0 \int_{-R}^R (z-a) \left(\frac{1}{\sqrt{R^2 - 2az + a^2}} - \frac{1}{a-z} \right) dz$$

(A)

分部积分

$$\bullet \Rightarrow -2\pi k m_0 \rho_0 \left(\int_R^R \frac{(z-a) dz}{\sqrt{R^2 - 2az + a^2}} + 2R \right)$$

$$-2\pi k m_0 \rho_0 \left[\left(-\frac{1}{a}\right) \int_R^R (z-a) d\sqrt{R^2 - 2az + a^2} + 2R \right]$$

$$= -2\pi k m_0 \rho_0 \left(\frac{2R^3}{3a^2} - 2R + 2R \right) = -\frac{\left(\frac{4}{3} 2R^3 \rho_0\right) m_0 k}{a^2} = -k \frac{M_0 m_0}{a^2}$$

$$M_0 \triangleq \frac{4}{3} 2R^3 \rho_0 \Rightarrow \vec{F} = \left(0, 0, -\frac{k m_0 M_0}{a^2} \right)$$

例6. 求 $\Omega = \{(x,y) \mid | \frac{x+y}{\sqrt{2}} - x^2 y^2 | \leq 1\}$ 的面积

由曲线围成得 Ω 的面积 $V(\Omega) = \iint_D \left| \frac{x+y}{\sqrt{2}} - x^2 y^2 \right| d\sigma$ (ch10 例4)

解: 设 $D_1: (x-\frac{1}{\sqrt{2}})^2 + (y-\frac{1}{\sqrt{2}})^2 \leq (\frac{1}{2})^2$, $D_2 = D - D_1$ 则 $D = D_1 + D_2$

$$V(\Omega) = \iint_D \left| \frac{x+y}{\sqrt{2}} - x^2 y^2 \right| d\sigma = \iint_{D_1} \left| \frac{x+y}{\sqrt{2}} - x^2 y^2 \right| d\sigma + \iint_{D_2} \left| \frac{x+y}{\sqrt{2}} - x^2 y^2 \right| d\sigma$$

$$\bullet \text{ 令 } g(x,y) = \frac{x+y}{\sqrt{2}} - x^2 y^2 \text{ 则 } V(\Omega) = \iint_{D_1} g(x,y) d\sigma - \iint_{D_2} g(x,y) d\sigma$$

$$= \iint_{D_1} g(x,y) d\sigma - \iint_{D_1 + D_2} g(x,y) d\sigma \quad \checkmark \text{ 证}$$


$$= \iint_{D_1} g(x,y) d\sigma \frac{x-\frac{1}{\sqrt{2}} = r \cos \theta}{y-\frac{1}{\sqrt{2}} = r \sin \theta} = \int_0^{2\pi} \int_0^{\frac{1}{2}} \left(\frac{1}{\sqrt{2}} + r^2 \right) r dr d\theta = \frac{\pi}{16}$$

$$\iint_{D_1 + D_2} g(x,y) d\sigma = \iint_{x^2 + y^2 \leq 1} \left(\frac{x+y}{\sqrt{2}} - x^2 y^2 \right) d\sigma = 0 - \iint_{x^2 + y^2 \leq 1} x^2 y^2 d\sigma = -\frac{\pi}{2}$$

$$\therefore V(\Omega) = \frac{\pi}{16} - \left(-\frac{\pi}{2}\right) = \frac{9}{16}\pi$$

(5)

例7. 设两根长为 a , 质量为 m_0 的均匀细杆位于同一直线上, 其左端距离为 a , 求两杆间万有引力 F .



解: 设两细杆所在直线为 Ox 轴. 两杆的线密度均为 $\rho_0 = \frac{m_0}{a}$

$$F = \int_{2a}^{3a} \int_0^a \frac{k \left(\frac{m}{a} dx \right) \left(\frac{m}{a} dx_0 \right)}{(x_0 - x)^2} = \left(\frac{m}{a} \right)^2 k \int_{2a}^{3a} \left(\int_0^a \frac{dx}{(x_0 - x)^2} \right) dx_0$$

$$= k \left(\frac{m}{a} \right)^2 \int_{2a}^{3a} \left(\frac{1}{x_0 - a} - \frac{1}{x_0} \right) dx_0 = k \left(\frac{m}{a} \right)^2 \ln \frac{x_0 - a}{x_0} \Big|_{2a}^{3a} = k \left(\frac{m}{a} \right)^2 \ln \frac{4}{3}$$

(二). 作业: P110.3: $5/8$; 6; 12; 14; 16; 19; Ch10 总/4.

(三). 第24讲: n 重积分: $\iint \dots \int_{\Omega_n} f(x_1, x_2, \dots, x_n) dx_1 dx_2 \dots dx_n$

其中, Ω_n 是 n 维空间 \mathbb{R}^n 中的有界闭区域, $f(x_1, x_2, \dots, x_n)$

在 Ω_n 中通常连续. 至少是确定且有限.

(6).